Home Care Schedule (2008.7.15) Schedule Main Operation Support Operation 1,Spurt the medicinal powder with saline into Trachesomy tube 2, Perform left-bodyflip and back-rap · remove sputum 3. Disinfect tracheostomy 4. Inspect humidifier and keep water level above red line and keep knob setting at "6" for humidity level 5. 06:00-06:45 Massage the belly clockwise around large intestine for 15min. 6. Changes dirty clothes Disinfect urethra mouth and anus - apply iodine and saline or other instructed medicne, and change Y gauze on tracheostomy 8. Remove residue water in tubes and collector etc. 9. Measure body temperature and blood oxygen %, and glucose level 06:45-07:00 1. Apply medicine for before meal 07:30-08:00: Provide drinking water, wash 07:00-07:45 Feed a spoon of milk powder with water 2.Break for own breakfast urine bottle, take out dirty clothes, follow up to

1. Apply medicine for after meal 2. Apply liquid medicines as instructed

07:45-08:00

15:00-16:00

16:00-17:00

17:00-17:30

17:30-18:00

18:00-18:45

18:45-19:00

19:00-19:45

19:45-20:00

20:00-22:00

22:00-23:00

23:00-23:30

23:30-

03:00(02:00) 02:00:00 AM

(overcast)

02:30-03:00 (fair-

weather)

04:00:00 AM

(overcast) 03:00-06:00

Attention

Break for own nap

Break for own bath

Take a sponge bath

1.Apply medicine for before meal

knob setting at "7" for humidity level

Break for night sleep

assist for break time

2. Perform right-body-flip

and back-rap · remove sputum 3. Inspect

Assist to wheelchair 1. Feed drinking water 2. Exercise Body

19:30: Follow up to

21:00 Provide drinking

02:30 Measure Vital Sign

and double check all the

equipement

water, clean up urine

bottle, change dirty

assist break time

movement 3. Assist to

humidifer

bed

clothes

| | Massage the belly around large intestine 4. Inspection humified and water level 5. Brush teeth 6. Remove residue water in tubes and collectors etc. | |
|-------------|--|---|
| 9:00-10:00 | 1. Sit to wheelchair and feed two spoons of milk with water 2. Feed drinking water | 09:00: Assist to wheelchair and feed milk 09:15: Assit to bed |
| 10:00-10:20 | Exercise hands and feet 2. Massage the belly 3. Exercise other body movement | Assist to wheelchair, change dirty bed sheet |
| 10:20-11:00 | Move to bed 2. Exercise body movement (foot and hand) | 11:00: Assist to bed |
| 11:00-11:45 | Perform left-body-flip and back-rap - remove sputum 2. Disinfect tracheostomy 3. Inspect humidifier 4. Change Y gauze on tracheostomy | |
| 11:45-12:00 | Apply medicine for before meal | |
| 12:00-12:45 | Feed two spoons of milk powder with drinking water Move to wheelchair Apply medicine for after meal Measure body temperature and blood oxygen | Provide drinking water, ibid as main operation |
| 12:45-13:00 | | Follow up to assist break time 2. Apply medicine for after meal |

1. Perform right-body-flip and back-rap · remove sputum 2. Disinfect tracheostomy 3.

| 11:00-11:45 | Perform left-body-flip and back-rap - remove sputum 2. Disinfect tracheostomy 3. Inspect humidifier 4. Change Y gauze on tracheostomy | |
|-------------|---|---|
| 11:45-12:00 | Apply medicine for before meal | |
| 12:00-12:45 | | Provide drinking water, ibid as main operation |
| 12:45-13:00 | | Follow up to assist break time 2. Apply medicine for after meal |
| 13:00-15:00 | | 1. Follow up to assist break time 2. 14:45: Feed nutrient |
| | | 1. Apply liquid medicines |

Feed two spoons of milk power with drinking water 2. Move to wheelchair

and back-rap - remove sputum = 3. Inpsect humidifier=4. Change Y gauze on

1. Apply medicine for after meal 2. Apply liquid medicines as instructed

tracheostomy 4. Disinfect uretha mouth and anus

Measure body temperature and blood oxygen %

humidifier and keep water level above red line

Complete putum suction before feeding milk is a must

4. Do not let fluid medicine leaked into tracheostomy

Spurt medicinal powder with saline into tracheostomy tube 2. Perform left-body-flip

1. Feed two spoons of milk powder with drinking water 2, 19:30: Break for own dinner

Perform right-body-flip and back-rap · remove sputum and disinfect tracheostomy

connector area 2. Replace Y gauze of Tracheostomy 3. Disinfect uretha mouth and

1.Feed a spoon of milk powder 2. Fix urine bag 3. Put towel underneath clothing 4.

Perform right-body-flip and back-rap · remove sputum 2. Feed drinking water

Perform left-body-flip and back-rap · remove sputum 2. Feed drinking water

1. Own break for sleep but alert and action to alarm and abnormal condition

5. Perform feeding milk 15-30 minutes after applied medicine before meal Before breaking for sleep need to check all the connector are set up properly

1. Perform right-body-flip and back-rap · remove sputum 2. Feed drinking water 3.

Change urine bag 4. Apply iodine and saline solution where-ever needed 5. Inspect

3. Need to shake the medicine bottle before spurting the medicine into tracheostomy

1. Sequence of Putum Suction @ tracheostomy : power on suction machine, wear decinfected glove, connect O2

to pressure-balloon and press 8-time, if putum too thick and drop 3 droplets of saline water into tracheostomy,

1. Perform left-body-flip and back-rap · remove sputum 2. Inspect humidifer and keep